

Existence of Periodic Solutions of Rayleigh Equations with Singularity

Shile Zhou, Shiping Lu

*School of Math and Statistics, Nanjing University of Information Science & Technology,
Nanjing 210044, China*

(Received March 21 2019, accepted September 20 2019)

Abstract: The existence of positive periodic solutions is studied for the Rayleigh equation with a singularity of attractive type

$$x''(t) + f(x'(t)) - \varphi(t)x(t) + \frac{\alpha(t)}{x^\mu} = h(t),$$

Where $f: R \rightarrow R$ is continuous function, which has a singularity at $x = 0$, that is $\lim_{x \rightarrow 0} \frac{\alpha(t)}{x^\mu} = \infty$. A new result on the existence of T-periodic positive solution to the above equation is obtained. The methods are based on Mawhin's continuation theorem of coincidence degree theory.

Keywords: Differential Equation; Singularity; Periodic Solution; Mawhin's Continuation Theorem, Coincidence Degree Theory.

1. Introduction

Second order differential equations are often used to model some dynamic problems in real world, where the problems are related to the first derivative with respect to the state variable, such as the resistance of air to a body in free falling movement, the resistance of moving charges in liquid, etc. Differential equation with a singularity is derived from physics, biology and other fields. For example, the famous Rayleigh-plesset equation[1-2]

$$u''(t) - \frac{4\mu}{u^{\frac{4}{5}}(t)}u'(t) + \varphi(t)u^{\frac{1}{5}}(t) + \frac{A}{u^{\frac{6k-1}{5}}(t)} - \frac{5S}{u^{\frac{1}{5}}(t)} = 0$$

describes the oscillation of spherical bubbles affected by periodic sound fields in a liquid. The Rayleigh-plesset equation plays an important role in fluid dynamics. It can be derived by taking spherical coordinates in the Euler equation and assuming some physically acceptable simplifications. Many physical, biological, and medical models related to cavitations and luminescence depend on this equation.

The study of differential equations with singularities began with the publication of Nagumo [3], and the interest in this field increased with the paper [4] of Lazer and Solimini. In that paper, the periodic problem of equations as follows

$$u''(t) - \frac{1}{u^{\alpha(t)}} = h(t) \text{ (repulsive type)} \quad (1.1)$$

and

$$u''(t) + \frac{1}{u^{\alpha(t)}} = h(t) \text{ (attractive type)} \quad (1.2)$$

was investigated. Inspired by this, later, many scholars devoted themselves to the study of the problems of existence and stability of periodic solutions for singular equations [5-9]. Among these papers, most of them considered the problem of periodic solutions for differential equations with repulsive singularities. The methods are mainly based on the theory of coincidence degree established by Mawhin [10], which has become one of the key methods to study the existence of periodic solutions for differential equations of many types. For example, Lu and Kong studied the periodic solution problem for mean curvature equation[8]

$$\left(\frac{u'(t)}{\sqrt{1 + (u'(t))^2}} \right)' + f(u(t))u'(t) + g(u(t - \sigma)) = e(t)$$

where $0 \leq \sigma \leq T$, $g: (0, +\infty) \rightarrow R$ is a continuous function, and $g(x)$ has a singularity at $x = 0$.

The behaviors of Rayleigh equation have been widely investigated due to their applications in many fields such as physics, mechanics and the engineering technique fields [11-19]. The authors in [14] studied the existence of periodic positive solutions for a Rayleigh equation with a singularity of attractive type

$$x''(t) - f(x'(t)) + g(t, x) = 0. \quad (1.3)$$

By using a continuation theorem of coincidence degree principle, they obtained the following result.

Theorem 1.1 Suppose $f(0) < 0$ and the following conditions are satisfied:

1. $\lim_{x \rightarrow 0^+} \inf_{t \in [0, T]} g(t, x) = +\infty$;
2. There is a constant $M > 0$ for any $(t, u) \in [0, T] \times (M, \infty)$, the relation $g(t, u) < -f(0)$ holds.

Then, Rayleigh equation (1.3) has at least one T -periodic positive solution.

Later, Guo and others made a further research [17], in which they studied the existence of periodic solutions of Rayleigh equation with a singularity of repulsive type

$$x''(t) + f(x'(t)) + \varphi(t)x(t) - \frac{1}{x^\alpha(t)} = p(t),$$

where $f: \mathbb{R} \rightarrow \mathbb{R}$ is continuous, $f(0) = 0$, and $\alpha \geq 1$ is a constant, $\varphi, p: \mathbb{R} \rightarrow \mathbb{R}$ are continuous with T -periodic with respect to variable t .

The aim of this paper is to study the periodic problem for the Rayleigh equation with a singularity of attractive type.

$$x''(t) + f(x'(t)) - \varphi(t)x(t) + \frac{\alpha(t)}{x^\mu(t)} = h(t), \quad (1.4)$$

where $f: \mathbb{R} \rightarrow \mathbb{R}$ is continuous, $f(0) = 0$, and φ, α, h are T -periodic continuous functions, $\max_{t \in [0, T]} h(t) > 0$,

$\bar{\varphi} < 0, \alpha(t) > 0$ for $t \in [0, T], \mu > 0$ is constant. By means of a continuation theorem of coincidence degree theory, a new result on the existence of periodic solutions to equation (1.4) is obtained. The interesting is that the sign of $\varphi(t)$ is allowed to change. In such a situation, the priori upper bound of periodic solutions associated with equation (1.4) is more difficult to estimate than the case of $\varphi(t) \geq 0$ for $t \in [0, T]$. In order to overcome this difficulty, the priori upper bound of periodic solutions to equation (1.4) is obtained by using extreme value principle. This method is essentially different to the corresponding ones used by the known literature.

2. Preliminary lemmas

In this section, let $C_T := \{x \in C(\mathbb{R}, \mathbb{R}), x(t+T) = x(t), \forall t \in \mathbb{R}\}$ with the norm $\|x\|_\infty = \max_{t \in [0, T]} |x(t)|$, $C_T^1 := \{x \in C^1(\mathbb{R}, \mathbb{R}), x(t+T) = x(t), \forall t \in \mathbb{R}\}$ with the norm $\|x\|_{C_T^1} := \max\{\|x\|_\infty, \|x'\|_\infty\}$. Clearly, C_T and C_T^1 are both Banach spaces. For any $y \in C_T$, let $y_+(t) := \max\{y(t), 0\}$, $y_-(t) := -\max\{y(t), 0\}$, $\bar{y} = \frac{1}{T} \int_0^T y(s) ds$, $y_M = \max_{t \in [0, T]} y(t)$, $y_m = \min_{t \in [0, T]} y(t)$, $\|y\|_p := (\int_0^T |y(s)|^p ds)^{\frac{1}{p}}$, $p \in [1, +\infty)$. Clearly, $y(t) = y_+(t) - y_-(t)$, $\bar{\varphi} = \bar{\varphi}_+ - \bar{\varphi}_-$ for all $t \in \mathbb{R}$.

Lemma 2.1 Assume that there exist positive constants of N_0, N_1 and N_2 with $N_0 < N_1$ such that the following conditions hold.

[C1] For each $\lambda \in (0, 1]$, each possible positive T -periodic solution x to the equation

$$u''(t) + \lambda f(u'(t)) - \lambda \varphi(t)u(t) + \frac{\lambda \alpha(t)}{u^\mu(t)} = \lambda h(t)$$

satisfies the inequalities $N_0 < u(t) < N_1$ and $|u'(t)| < N_2$ for all $t \in [0, T]$.

[C2] Each possible solution c to the equation

$$\frac{\bar{\alpha}}{c^\mu} - c\bar{\varphi} - \bar{h} = 0$$

satisfies the inequality $N_0 < c < N_1$.

[C3] The inequality

$$(\frac{\bar{\alpha}}{N_0^\mu} - N_0\bar{\varphi} - \bar{h})(\frac{\bar{\alpha}}{N_1^\mu} - N_1\bar{\varphi} - \bar{h}) < 0$$

holds.

Then equations (1.4) has at least one positive T -periodic solution u such that $N_0 \leq u(t) \leq N_1$ and $|u'(t)| \leq N_2$ for all $t \in [0, T]$.